



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Multiethnic Exome-Wide Association Study of Subclinical Atherosclerosis

Citation for published version:

CHARGE Consortium, Natarajan, P, Bis, JC, Bielak, LF, Cox, AJ, Dörr, M, Feitosa, MF, Franceschini, N, Guo, X, Hwang, S-J, Isaacs, A, Jhun, MA, Kavousi, M, Li-Gao, R, Lyytikäinen, L-P, Marioni, RE, Schminke, U, Stitzel, NO, Tada, H, van Setten, J, Smith, AV, Vojinovic, D, Yanek, LR, Yao, J, Yerges-Armstrong, LM, Amin, N, Baber, U, Borecki, IB, Carr, JJ, Chen, Y-DI, Cupples, LA, de Jong, PA, de Koning, H, de Vos, BD, Demirkan, A, Fuster, V, Franco, OH, Goodarzi, MO, Harris, TB, Heckbert, SR, Heiss, G, Hoffmann, U, Hofman, A, Işgum, I, Jukema, JW, Kähönen, M, Kardia, SLR, Kral, BG, Wardlaw, JM, Deary, IJ & Wilson, JG 2016, 'Multiethnic Exome-Wide Association Study of Subclinical Atherosclerosis', *Circulation: Cardiovascular Genetics*. <https://doi.org/10.1161/CIRCGENETICS.116.001572>

Digital Object Identifier (DOI):

[10.1161/CIRCGENETICS.116.001572](https://doi.org/10.1161/CIRCGENETICS.116.001572)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Circulation: Cardiovascular Genetics

Publisher Rights Statement:

This is author's peer-reviewed manuscript as accepted for publication

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



1 Multiethnic Exome-Wide Association Study of Subclinical Atherosclerosis

2 Pradeep Natarajan MD MMSc^{1,2*}, Joshua C. Bis PhD^{3*}, Lawrence F. Bielak DDS MPH⁴,
 3 Amanda J. Cox PhD⁵, Marcus Dörr MD^{6,7}, Mary F. Feitosa PhD⁸, Nora Franceschini MD MPH⁹,
 4 Xiuqing Guo PhD¹⁰, Shih-Jen Hwang PhD¹¹, Aaron Isaacs PhD^{12,13}, Min A. Jhun PhD⁴, Maryam
 5 Kavousi MD PhD¹², Ruifang Li-Gao MSc¹⁴, Leo-Pekka Lyytikäinen MD^{15,16}, Riccardo E.
 6 Marioni PhD^{17,18,19}, Ulf Schminke MD²⁰, Nathan O. Stitziel MD PhD^{8,21,22}, Hayato Tada MD²³,
 7 Jessica van Setten PhD²⁴, Albert V. Smith PhD^{25,26}, Dina Vojinovic MD MSc¹², Lisa R. Yanek
 8 MPH²⁷, Jie Yao MD MS¹⁰, Laura M. Yerges-Armstrong PhD²⁸, Najaf Amin PhD¹², Usman
 9 Baber MD²⁹, Ingrid B. Borecki PhD³⁰, J. Jeffrey Carr MD MSc³¹, Yii-Der Ida Chen PhD¹⁰, L.
 10 Adrienne Cupples PhD³², Pim A. de Jong MD PhD³³, Harry de Koning PhD³⁴, Bob D. de Vos
 11 MSc³⁵, Ayse Demirkan PhD¹², Valentin Fuster MD PhD²⁹, Oscar H. Franco MD PhD¹², Mark O.
 12 Goodarzi MD PhD³⁶, Tamara B. Harris MD³⁷, Susan R. Heckbert MD PhD³⁸, Gerardo Heiss MD
 13 PhD⁹, Udo Hoffmann MD PhD³⁹, Albert Hofman MD PhD¹², Ivana Išgum PhD³⁵, J. Wouter
 14 Jukema MD PhD^{40,41,42}, Mika Kähönen MD PhD^{43,44}, Sharon L.R. Kardia PhD⁴, Brian G. Kral
 15 MD MPH²⁷, Lenore J. Launer PhD³⁷, Joe Massaro PhD³², Roxana Mehran MD²⁹, Braxton D.
 16 Mitchell PhD MPH^{28,45}, Thomas H. Mosley Jr. PhD⁴⁶, Renée de Mutsert PhD¹⁴, Anne B.
 17 Newman MD⁴⁷, Khanh-dung Nguyen PhD⁴⁸, Kari E. North PhD⁹, Jeffrey R. O'Connell PhD²⁸,
 18 Matthijs Oudkerk MD⁴⁹, James S. Pankow PhD MPH⁵⁰, Gina M. Peloso PhD^{1,2,32}, Wendy Post
 19 MD MS²⁷, Michael A. Province PhD⁸, Laura M. Raffield PhD⁵, Olli T. Raitakari MD PhD^{51,52},
 20 Dermot F. Reilly PhD⁴⁸, Fernando Rivadeneira MD PhD^{12,53,54}, Frits Rosendaal MD PhD¹⁴,
 21 Samantha Sartori PhD²⁹, Kent D. Taylor PhD¹⁰, Alexander Teumer PhD^{7,55}, Stella Trompet
 22 PhD⁴⁰, Stephen T. Turner MD⁵⁶, Andre G. Uitterlinden PhD^{12,53,54}, Dhananjay Vaidya PhD MPH
 23 MBBS²⁷, Aad van der Lugt MD PhD⁵⁷, Uwe Völker PhD^{7,58}, Joanna M. Wardlaw MD^{17,59},
 24 Christina L. Wassel PhD MS⁶⁰, Stefan Weiss PhD^{7,58}, Mary K. Wojczynski PhD⁸, Diane M.
 25 Becker ScD PhD²⁷, Lewis C. Becker MD²⁷, Eric Boerwinkle PhD^{61,62}, Donald W. Bowden PhD⁵,
 26 Ian J. Deary PhD^{17,63}, Abbas Dehghan MD PhD¹², Stephan B. Felix MD^{6,7}, Vilmundur Gudnason
 27 MD PhD^{25,26}, Terho Lehtimäki MD PhD^{15,16}, Rasika Mathias ScD²⁷, Dennis O. Mook-Kanamori
 28 MD PhD^{14,64}, Bruce M. Psaty MD^{3,38,65,66}, Daniel J. Rader MD⁶⁷, Jerome I. Rotter MD¹⁰, James
 29 G. Wilson MD⁶⁸, Cornelia M. van Duijn PhD^{12,54}, Henry Völzke MD^{7,55,69}, Sekar Kathiresan
 30 MD^{1,2}, Patricia A. Peyser PhD⁴, Christopher J. O'Donnell MD MPH^{70†}, CHARGE Consortium

32 * Contributed equally

33 † Corresponding author

35 ¹ Center for Human Genetic Research and Cardiovascular Research Center, Massachusetts General Hospital,
 36 Boston, MA, 02114

37 ² Program in Medical and Population Genetics, Broad Institute of Harvard and MIT, Cambridge, MA 02142

38 ³ Cardiovascular Health Research Unit, Department of Medicine, University of Washington, Seattle, WA 98101

39 ⁴ Department of Epidemiology, School of Public Health, University of Michigan, Ann Arbor, MI 48109

40 ⁵ Diabetes Heart Study, Wake Forest Health Sciences, Winston-Salem, NC 27104

41 ⁶ Department of Internal Medicine, University Medicine Greifswald, Greifswald, Germany 17475

42 ⁷ DZHK (German Centre for Cardiovascular Research), partner Site Greifswald, Germany 17475

43 ⁸ Division of Statistical Genetics, Department of Genetics, Washington University School of Medicine, St. Louis,
 44 MO 63108

45 ⁹ Epidemiology, Gillings School of Global Public Health, University of North Carolina, Chapel Hill, NC 27514

46 ¹⁰ Institute for Translational Genomics and Population Sciences, Los Angeles Biomedical Research Institute, and
 47 Department of Pediatrics at Harbor-UCLA Medical Center, Torrance, CA 90502

- 48 ¹¹ Framingham Heart Study, National Heart Lung and Blood Institute / National Institutes of Health, Framingham ,
49 MA 01701
- 50 ¹² Genetic Epidemiology Unit, Department of Epidemiology, Erasmus MC, University Medical Center Rotterdam,
51 Rotterdam, NL
- 52 ¹³ CARIM School for Cardiovascular Diseases, Maastricht Centre for Systems Biology, Department of
53 Biochemistry, Maastricht University, Maastricht, NL
- 54 ¹⁴ Department of Clinical Epidemiology, Leiden University Medical Center, Leiden, NL
- 55 ¹⁵ Department of Clinical Chemistry, Fimlab Laboratories, Tampere, Finland 33520
- 56 ¹⁶ Department of Clinical Chemistry, University of Tampere School of Medicine, Finland, 33014
- 57 ¹⁷ Centre for Cognitive Ageing and Cognitive Epidemiology, University of Edinburgh, Edinburgh, UK, EH8 9JZ
- 58 ¹⁸ Centre for Genomic and Experimental Medicine, Institute of Genetics and Molecular Medicine, University of
59 Edinburgh, Edinburgh, UK, EH4 2XU
- 60 ¹⁹ Queensland Brain Institute, University of Queensland, Brisbane, QLD, Australia 4072
- 61 ²⁰ Department of Neurology, University Medicine Greifswald, Greifswald, Germany 17475
- 62 ²¹ Cardiovascular Division, Department of Medicine, Washington University School of Medicine, St. Louis, MO
63 63110
- 64 ²² McDonnell Genome Institute, Washington University School of Medicine, St. Louis, MO 63110
- 65 ²³ Division of Cardiovascular Medicine, Kanazawa University Graduate School of Medicine, Kanazawa, Japan
66 9208641
- 67 ²⁴ Cardiology, University Medical Center Utrecht, Utrecht, NL
- 68 ²⁵ Icelandic Heart Association, Kopavogur, Iceland
- 69 ²⁶ Faculty of Medicine, University of Iceland, Reykjavik, Iceland
- 70 ²⁷ Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, MD 21287
- 71 ²⁸ Division of Endocrinology, Diabetes & Nutrition, Department of Medicine, University of Maryland School of
72 Medicine, Baltimore, MD 21201
- 73 ²⁹ Cardiovascular Institute, Icahn School of Medicine at Mount Sinai, New York, NY 10029
- 74 ³⁰ Analytical Genetics, Regeneron Pharmaceuticals Inc, Tarrytown, NY 10591
- 75 ³¹ Radiology, Cardiovascular Medicine and Biomedical Informatics, Vanderbilt, Nashville, TN 37203
- 76 ³² Department of Biostatistics, Boston University School of Public Health, Boston, MA 02118
- 77 ³³ Department of Radiology, University Medical Center Utrecht, Utrecht, NL
- 78 ³⁴ Department of Public Health, Erasmus Medical Center Rotterdam, Rotterdam, NL
- 79 ³⁵ Image Sciences Institute, University Medical Center Utrecht, Utrecht, NL
- 80 ³⁶ Division of Endocrinology, Diabetes and Metabolism, Cedars-Sinai Medical Center, Los Angeles, CA 90048
- 81 ³⁷ Laboratory of Epidemiology and Population Sciences, National Institute on Aging Intramural Research / National
82 Institutes of Health, Baltimore, MD 20892
- 83 ³⁸ Department of Epidemiology, University of Washington, Seattle, WA 98101
- 84 ³⁹ Department of Radiology, Massachusetts General Hospital, Boston, MA 02114
- 85 ⁴⁰ Department of Cardiology, Leiden University Medical Center, Leiden, NL
- 86 ⁴¹ Durrer Center for Cardiogenetic Research, Netherlands Heart Institute, Amsterdam, NL
- 87 ⁴² Interuniversity Cardiology Institute of the Netherlands, Netherlands Heart Institute, Utrecht, NL
- 88 ⁴³ Department of Clinical Physiology, Tampere University Hospital, Tampere, Finland 33521
- 89 ⁴⁴ Department of Clinical Physiology, University of Tampere School of Medicine, Tampere, Finland, 33014
- 90 ⁴⁵ Geriatrics Research and Education Clinical Center, Baltimore Veterans Administration Medical Center,
91 Baltimore, MD 21201⁴⁶ Division of Geriatric Medicine, Department of Medicine, University of Mississippi Medical
92 Center, Jackson, MS 39216
- 93 ⁴⁷ Department of Epidemiology, Graduate School of Public Health, University of Pittsburgh, Pittsburgh, PA 15261
- 94 ⁴⁸ Genetics & Pharmacogenomics Department, Merck Sharpe & Dohme Corp., Boston, MA 02115
- 95 ⁴⁹ Center for Medical Imaging - North East Netherlands, University Medical Center Groningen, University of
96 Groningen, Groningen, NL
- 97 ⁵⁰ Division of Epidemiology and Community Health, University of Minnesota, Minneapolis, MN 5545
- 98 ⁵¹ Department of Clinical Physiology and Nuclear Medicine, Turku University Hospital, Turku, Finland 20521
- 99 ⁵² Research Centre of Applied and Preventive Cardiovascular Medicine, University of Turku, Turku, Finland 20520
- 100 ⁵³ Department of Internal Medicine, Erasmus MC, University Medical Center Rotterdam, Rotterdam, NL
- 101 ⁵⁴ Netherlands Consortium for Healthy Aging, Rotterdam, NL
- 102 ⁵⁵ Institute for Community Medicine, University Medicine Greifswald, Greifswald, Germany 17475⁵⁶ Division of
103 Nephrology and Hypertension, Department of Internal Medicine, Mayo Clinic, Rochester, MN 55905

- ⁵⁷ Department of Radiology, Erasmus MC, University Medical Center Rotterdam, Rotterdam, NL
⁵⁸ Interfaculty Institute of Genetics and Functional Genomics, University Medicine Greifswald, Greifswald, Germany, 17475
⁵⁹ Division of Neuroimaging Sciences and Brain Research Imaging Centre, Centre for Clinical Brain Sciences, University of Edinburgh, Edinburgh, UK, EH4 2XU
⁶⁰ Department of Pathology and Laboratory Medicine, The University of Vermont, Colchester, VT 05446
⁶¹ Human Genetics Center, The University of Texas Health Science Center at Houston, Houston, TX 77030
⁶² Epidemiology, School of Public Health, The University of Texas Health Science Center at Houston, Houston, TX 77030
⁶³ Department of Psychology, University of Edinburgh, Edinburgh, UK EH8 9JZ
⁶⁴ Department of Public Health and Primary Care, Leiden University Medical Center, Leiden, NL
⁶⁵ Department of Health Services, University of Washington, Seattle, WA 98101
⁶⁶ Group Health Research Institute, Group Health Cooperative, Seattle, WA 98101
⁶⁷ Division of Translational Medicine & Human Genetics, Department of Medicine, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104
⁶⁸ Department of Physiology and Biophysics, University of Mississippi Medical Center, Jackson, MS 39216
⁶⁹ German Center for Diabetes Research (DZD), Greifswald, Germany 17475
⁷⁰ Division of Cardiology, Boston Veterans Affairs Healthcare System, Boston, MA 02132

Word count (abstract): 243

Word count (abstract + text + legends + tables + disclosures + funding + references): 5,639

References: 44

Running title: Exome association with subclinical atherosclerosis

Please address correspondence to:

Christopher J. O'Donnell, MD MPH

Associate Professor Medicine, Harvard Medical School

Chief of Cardiology, Boston Veterans Affairs Healthcare System

1400 Veterans of Foreign Wars Parkway, Building 1 5B-113

Boston, MA 02132 USA

Phone: 857-203-6840, Fax: 857-203-5550

Email: odonnellc@nhlbi.nih.gov

Abstract

Background – The burden of subclinical atherosclerosis in asymptomatic individuals is heritable and associated with elevated risk of developing clinical coronary heart disease (CHD). We sought to identify genetic variants in protein-coding regions associated with subclinical atherosclerosis and the risk of subsequent CHD.

Methods and Results – We studied a total of 25,109 European ancestry and African-American participants with coronary artery calcification (CAC) measured by cardiac computed tomography and 52,869 with common carotid intima media thickness (CIMT) measured by ultrasonography within the Cohorts for Heart and Aging Research in Genomic Epidemiology (CHARGE) Consortium. Participants were genotyped for 247,870 DNA sequence variants (231,539 in exons) across the genome. A meta-analysis of exome-wide association studies was performed across cohorts for CAC and CIMT. *APOB* p.Arg3527Gln was associated with four-fold excess CAC ($P = 3 \times 10^{-10}$). The *APOE* $\epsilon 2$ allele (p.Arg176Cys) was associated with both 22.3% reduced CAC ($P = 1 \times 10^{-12}$) and 1.4% reduced CIMT ($P = 4 \times 10^{-14}$) in carriers compared with non-carriers. In secondary analyses conditioning on LDL cholesterol concentration, the $\epsilon 2$ protective association with CAC, although attenuated, remained strongly significant. Additionally, the presence of $\epsilon 2$ was associated with reduced risk for CHD (OR 0.77; $P = 1 \times 10^{-11}$).

Conclusions – Exome-wide association meta-analysis demonstrates that protein-coding variants in *APOB* and *APOE* associate with subclinical atherosclerosis. *APOE* $\epsilon 2$ represents the first significant association for multiple subclinical atherosclerosis traits across multiple ethnicities as well as clinical CHD.

Key Words: Genome Wide Association Study; exome; coronary artery calcification; carotid intima-media thickness; genomics

Background

Coronary heart disease (CHD) remains the leading cause of death and infirmity in developed countries.¹ Atherosclerosis is the underlying pathology of CHD.² The presence of atherosclerosis in individuals without clinical CHD, termed “subclinical atherosclerosis,” is associated with increased risk of developing clinical CHD independent of traditional risk factors prior to the onset of symptoms.³⁻⁶ Subclinical atherosclerosis is a heritable⁷⁻⁹ clinical phenotype that can be ascertained non-invasively as coronary artery calcification (CAC) by cardiac computed tomography (CT) and common carotid intima media thickness (CIMT) by carotid ultrasound.¹⁰

Genome-wide association studies (GWAS) within the Cohorts for Heart and Aging Research in Genomic Epidemiology (CHARGE) Consortium have discovered sites of common non-coding genetic variation associated with both CAC^{11, 12} and CIMT^{7, 11} among those of European ancestry. Non-coding single nucleotide polymorphisms (SNPs) at the 9p21 and 6p24 regions, near the *CDKN2A* and *PHACTR1* genes, respectively, are strongly associated with both CAC burden and myocardial infarction (MI).¹² The 8q24 (*ZHX2*), 19q13 (*APOC1*), and 8q23 (*PINX1*) loci are strongly associated with CIMT.⁷ Observed associations for subclinical atherosclerosis among individuals of European ancestry, however, have not been replicated in those of African ancestry.^{13, 14} Furthermore, since the biologic implications of non-coding variation are not as readily interpreted as with coding variation, the roles of such variants in human atherosclerosis remain unclear.¹⁵ Protein-coding variation tends to be infrequently observed and is often inadequately catalogued on earlier GWAS arrays.¹⁶ Rare genomic variation is not well-imputed and exome sequencing to detect such uncommon variation across large populations remains a costly endeavor. Here, we leverage the Illumina HumanExome BeadChip

array, enriched for protein-coding variation.¹⁷ We investigated whether there is evidence for associations of protein-coding variation with two measures of subclinical atherosclerosis across individuals of European and of African ancestry. And we further determine whether such DNA sequence variations may influence CHD risk.

Methods

Study Populations

The Illumina HumanExome Beadchip v1.0 or v1.1 (also known as the “exome chip”) was used to genotype participants across 19 cohorts of the CHARGE Consortium (**Supplement**).¹⁸ Participants with a diagnosis of CHD at the time of CAC phenotyping were excluded from CAC analysis. Participants who underwent carotid endarterectomy prior to CIMT phenotyping were excluded from CIMT analysis. 25,109 participants had CAC measured and 52,869 participants had CIMT measured. Each study received institutional review board approval, participants provided written informed consent, and respective governing ethics committees approved each study.

Measures

CAC Measurement

Cohorts used different CT scanners to ascertain CAC scoring (**Table S1**). CAC scoring by multidetector CT and by electron beam CT have been previously described to be highly concordant and are both recognized as valid tools to estimate CAC score.¹⁹⁻²¹ Total CAC score was quantified by the sum of CAC area weighted by density within individual coronary arteries by the Agatston method and the continuous score was used for analysis.²²

CIMT Measurement

Common carotid intima media thickness was derived by bilateral longitudinal common carotid artery analysis (imaging and measurement methods are described in the **Table S2**). The mean of the maximum thickness for each common carotid artery was the analytical variable.

Statistical Analyses

According to prespecified analysis plans, association analyses and meta-analyses were performed using the seqMeta package (<http://cran.r-project.org/web/packages/seqMeta/index.html>) in the R statistical software as has previously been performed for exome chip-based analyses.²³ To reduce skewness, CAC was natural log transformed after adding 1 and CIMT was natural log transformed. Each cohort performed an analysis for each genomic variant with the trait of interest independently and separately for individuals of European and African ancestry to minimize population biases. Covariates in the models included age, sex, and principal components of ancestry derived using EIGENSTRAT.²⁴ For studies with related samples, the pairwise kinship matrix was computed and accounted for in

the regression model. Score statistics and genotypic covariance matrices were computed for each cohort and used for additive single variant and gene-based analyses, respectively.

For our primary analyses, we tested the association of each genomic variant with CAC and with CIMT across all samples by meta-analysis that included all cohorts, irrespective of ancestry. We performed single variant analyses on variants that had a minor allele count of at least 20 and gene-based analyses for genes with combined minor allele frequency (MAF) of nonsynonymous variants at least 0.2% to reduce the likelihood of false positive results. We also performed two gene-based tests: 1) T1, where nonsynonymous variants with minor allele frequency (MAF) <1% were collapsed into a gene-based statistic, and 2) sequence kernel association test (SKAT) with MAF <5% for nonsynonymous variants to better account for collapsed variants with bidirectional phenotypic consequences. Regional association plots were generated using LocusZoom.²⁵ For our secondary analyses, we tested the association of each genomic variant with CAC and CIMT by meta-analysis separately among cohorts of European and African ancestry.

Given the 238,065 variants on the array that passed quality control, the Bonferroni-adjusted level of significance for single variant tests was $0.05/238,065 = 2.10 \times 10^{-7}$. Given the 17,574 genes with nonsynonymous variants on the array, the Bonferroni-adjusted level of significance for gene-based tests was $0.05/17,574 = 2.85 \times 10^{-6}$. For CAC, we had >90% power to detect a variant (MAF <1%) with effect size 0.31 standard deviations, or a gene (combined MAF <1%) with effect size 0.28 standard deviations at a sample size of 25,000. For CIMT, we had >90% power to detect a variant (MAF <1%) with effect size 0.21 standard deviations or a gene (combined

MAF <1%) with effect size 0.20 standard deviations with a sample size of 52,000. Power calculations were performed using the Genetic Power Calculator.²⁶

Methods for the secondary analyses are presented in the **Supplement**.

Results

Study Participants

19 cohorts participated in the meta-analyses of these two subclinical atherosclerotic traits and the clinical characteristics are summarized in **Table S1** and **Table S2**. A total of 25,109 participants were genotyped with the array and had CAC assessed; of these participants, 19,980 were of European ancestry and 5,129 were of African ancestry. 52,869 participants were genotyped and had CIMT assessed; 44,963 were of European ancestry and 7,906 were of African ancestry. 222,701 (93.5%) of the 238,065 variants were polymorphic in the CAC meta-analysis; of polymorphic variants, 193,373 (97.1%) were annotated as nonsynonymous or splice-site variants. Similarly, 227,344 (95.5%) of array variants were polymorphic in the CIMT meta-analysis and, of these, 217,235 (95.6%) were nonsynonymous or splice-site variants.

Coronary Artery Calcification Association

Figure 1 plots the meta-analysis CAC association P-value by genomic locus for each variant. The top loci with lead variants associated with CAC among all participants are listed in **Table 1**.

No systematic association inflation was observed across the set of statistical tests performed (Figure S1).

We identified previously-described common non-coding variant associations at the 9p21 and 6p24 loci. A 9p21 haplotype marked by lead SNP rs10757278-G (MAF 43%), an intergenic variant, was replicated and associated with increased CAC quantity (23.4%; 95% CI: 18.6, 28.3%; $P = 2 \times 10^{-24}$). Similarly, rs9349379-G (MAF 34%), an intronic variant within *PHACTR1*, was associated with increased CAC quantity (20.9%; 95% CI: 16.3, 25.8 %; $P = 5 \times 10^{-20}$). While these associations were robust for those of European ancestry, there was no apparent evidence for association in those of African ancestry (Figure S2, Figure S3). Both loci display locus heterogeneity, or multiple independent associations, for CAC in those of European ancestry (Table 1). We did not discover non-coding variants at other loci on the exome chip that met our stringent Bonferroni alpha threshold. Previously, rs3809346, an intronic variant of *COL4A2*, had a suggestive association with CAC,¹² but now in our European ancestry sample size that is twice as large, genome-wide significant association was not observed ($P = 2 \times 10^{-3}$).

Among functional variants, a nonsynonymous *APOB* (rs5742904-T; MAF 0.2%; NM_000384.2:c.10580G>A; NP_000375.2:p.Arg3527Gln) variant was significantly associated with CAC quantity. Carriers of the rare *APOB* missense variant had markedly increased CAC (4.1-fold; 95% CI: 2.6-, 6.4-fold; $P = 3 \times 10^{-10}$). In our meta-analysis, the Old Order Amish cohort primarily accounted for the strong association, and the variant was extremely rarely observed within other cohorts. Furthermore, the variant was not seen among individuals of African ancestry (Figure S4). We also discovered a distinct rare *APOB* missense variant (rs1801696-T;

MAF 0.6% European ancestry; NM_000384.2:c.7696G>A; NP_000375.2:p.Glu2566Lys), detected in individuals of European ancestry in most cohorts, that was moderately associated with increased CAC (1.9-fold; 95% CI: 1.6-, 2.1-fold; $P = 9 \times 10^{-6}$). This variant was not observed in individuals of African ancestry.

Additionally, a missense 19q13 variant within the *APOE* gene (rs7412-T; MAF 7.4% European ancestry, 10.8% African ancestry; NM_000041.2:c.526C>T; NP_000032.1:p.Arg176Cys) was associated with diminished CAC quantity (-22.3%; 95% CI, -27.6- -16.7%; $P = 1 \times 10^{-12}$) (**Figure S5**). This association was consistent in those of both European ancestry (-17.3%; 95% CI: -23.7, -10.3%; $P = 4 \times 10^{-6}$) and African ancestry (-35.2%; 95% CI: -43.6, -25.7%; $P = 5 \times 10^{-10}$) without significant heterogeneity ($P = 0.53$) (**Figure 2**). Additionally, an independent variant (rs769449-A; MAF 11% European ancestry, 2.4% African ancestry) within an intron of *APOE* also had nominal evidence of association with increased CAC quantity only in individuals of European ancestry (+15.0%; 95% CI: 7.9, 22.6%; $P = 2 \times 10^{-5}$).

To improve power of discovery for rare protein-coding variants, we conducted gene-based analyses by aggregating such variants within a gene into a single statistical unit to increase the exposure rate. However, collapsing nonsynonymous variants on the exome chip within a gene did not yield genome-wide significant results (**Figure S6**).

Carotid Intima Media Thickness Association

There was no systematic inflation of CIMT associations with any variant (**Figure S7**). The top meta-analysis association findings are listed in **Table 2** and a Manhattan plot of all associations is presented in **Figure 3**.

We noted that, in addition to diminished CAC, the rs7412-T *APOE* ϵ 2 allele was associated with diminished CIMT (-1.4%; 95% CI: -1.8, -1.0%; $P = 4 \times 10^{-14}$). There was consistency of association across European and African ancestry cohorts (**Figure 4** and **Figure S8**). There was no significant heterogeneity among the cohorts for this association (P heterogeneity = 0.23)

There were two additional independent suggestive associations at 19q13 at non-coding variants. A variant 5kb upstream of *LDLR* (rs11668477) was associated with diminished CIMT ($P = 5 \times 10^{-7}$) primarily among those of European ancestry. This variant has previously been associated with reduced LDL cholesterol.²⁷ The nearby rs7188-G variant (MAF 33% European ancestry, 7.9% African ancestry) within the 3'UTR region of *KANK2* was associated with CIMT in those of European ancestry ($P = 1 \times 10^{-6}$). Additionally, a rare missense variant (rs143873045-A; MAF 0.5% African ancestry; NM_001136191.2:c.1274C>T; NP_001129663.1:p.Ser425Leu) in *KANK2* only observed in individuals of African ancestry showed suggestive association with increased CIMT ($P = 4 \times 10^{-4}$). Lastly, in gene-based analyses, collapsing nonsynonymous variants within a gene did not yield significant associations (**Figure S9**).

***APOE* ϵ 2's Effect Conditional on LDL Cholesterol**

We sought to determine whether LDL cholesterol concentration accounted for the observed $\epsilon 2$ association with CAC. First, when restricting the original analysis only to participants with LDL cholesterol measurements ($n = 20,527$), $\epsilon 2$ remained significantly associated with reduced CAC quantity (-22.3%; 95% CI: -25.1, -19.3%; $P = 2 \times 10^{-11}$) (**Table S3**). When further adjusting for medication-adjusted LDL cholesterol, the effect estimate was diminished yet the association remained genome-wide significant (-17.0%; 95% CI: -19.7, -14.2%; $P = 2 \times 10^{-8}$).

APOE $\epsilon 2$'s Effect Conditional on $\epsilon 3$ and $\epsilon 4$

Given the absence of $\epsilon 4$ from the array, we sought to determine whether $\epsilon 2$'s apparent effect on reduced CAC quantity was due to a referent that includes a previously described risk allele ($\epsilon 3 + \epsilon 4$). 5,872 participants had CAC and the major *APOE* genotypes assessed by PCR. Each *APOE* genotype's association with CAC (to the $\epsilon 3/\epsilon 3$ referent) was performed by cohort and ethnicity and subsequently meta-analyzed with fixed effects. $\epsilon 2/\epsilon 3$ was associated with 10.8% reduced CAC (95% CI: -19.6, -0.01%; $P = 0.03$) and $\epsilon 2/\epsilon 2$ with 27.4 % reduced CAC (95% CI: -45.2, -0.04%; $P = 0.03$) (**Figure S10**).

Concordance of CHD Variants with Subclinical Atherosclerosis Associations

Of the 57 loci previously associated with CHD mainly in individuals of European or South Asian descent, 40 published variants were on the array and available for analysis. 32 of the 40 variants have the same effect direction for CAC and CHD ($P = 1.8 \times 10^{-4}$) whereas only 23 variants were concordant for CIMT ($P = 0.43$) in European ancestry participants (**Table S4**). When restricting

the analysis to variants with at least nominal association ($P < 0.05$) with CAC, all 17 had concordant effect directions ($P = 4.8 \times 10^{-7}$). A similar analysis with variants at least nominally associated with CIMT showed that 6 of 11 had concordant effect directions for CHD ($P = 0.56$).

Replication of Convergent Subclinical Atherosclerosis Finding with CHD

21,182 individuals of European ancestry, independent of the sample for subclinical atherosclerosis investigations, were genotyped by the Illumina HumanExome BeadChip array, of whom 9,472 had CHD.²⁸ In cross-sectional analyses, meta-analysis of rs7412-T confirmed a significantly lower odds of CHD (odds ratio 0.77; 95% CI: 0.71, 0.84; $P = 1.47 \times 10^{-10}$).

Discussion

In our exome-wide association analysis for subclinical atherosclerosis in two distinct ethnicities, we find that protein-coding mutations in *APOB* and *APOE* are associated with subclinical atherosclerosis. While the association for *APOB* was driven by a founder mutation in the Amish, a missense mutation in *APOE* ($\epsilon 2$) was associated with both reduced CAC and CIMT in individuals of European ancestry and African ancestry, even when adjusting for LDL cholesterol concentration. Furthermore, carriers of the $\epsilon 2$ allele had a reduced risk of coronary heart disease. Here, we provide evidence for the first exome-wide association across multiple subclinical atherosclerosis traits and multiple ethnicities for *APOE* $\epsilon 2$.

Both CAC and CIMT have been proposed as proximal clinical phenotypes of atherosclerosis that may identify individuals at high risk for developing clinical CHD. However, we see that alleles that associate with increased CHD risk also appear to largely result in increased CAC, which is less consistently observed with CIMT. This is concordant with the prior observation that CAC outperforms CIMT in predicting cardiovascular events.^{5, 29} Recently, *post hoc* analyses in statin trials to prevent cardiovascular disease observed that those with a higher burden of CHD-predisposing alleles are more likely to derive clinical benefit from preventive statin therapy.³⁰

The *APOB* p.Arg3527Gln (also known as p.Arg3500Gln) has been previously been shown to lead to increased concentrations of LDL cholesterol and premature CHD.³¹ Our association signal for this variant was nearly exclusively driven by the Old Order Amish, where it is known to be a founder mutation (MAF 12%) predisposing to increased LDL cholesterol concentrations and CAC quantity through disruption of the LDL receptor binding domain.³² We also observed a distinct *APOB* missense mutation, p.Glu2566Lys, with borderline association with increased CAC quantity. Unlike p.Arg3527Gln, p.Glu2566Lys does not occur within the LDL receptor binding domain but occurs within a conserved amphipathic motif of the β_2 domain predicted to influence the conversion of VLDL to LDL.³³

Furthermore, we demonstrated that *APOE* p.Arg176Cys (ϵ 2 allele) was associated with reduced CAC and reduced CIMT in both individuals of European and African ancestry. *APOE* is an essential mediator of the catabolism and clearance of triglyceride-rich and cholesterol-rich lipoproteins. The major alleles, ϵ 2, ϵ 3, and ϵ 4, have been previously linked to cardiovascular disease, from the candidate gene era, and ϵ 2 is the least common allele.^{34, 35} Previously, CHD

risk predisposition from $\epsilon 4$ was primarily thought to be mediated by LDL cholesterol raising effects but observations with $\epsilon 2$ have been mixed.³⁵ Similarly, $\epsilon 4$, unlike $\epsilon 2$, has been generally linked to ischemic stroke risk.³⁶ Major reasons for the lack of association of the major *APOE* alleles with cardiovascular traits in prior genome-wide association studies include the notable absence of rs7412 and rs429358 on population-based genotyping arrays as well as poor imputation of these variants. Similarly, rs429358 is not included on the array used for this study.

ApoE is a major ligand of LDL receptor and a key mediator of remnant lipoprotein particle clearance.^{37, 38} The $\epsilon 2$ allele is believed to result in less efficient LDL receptor binding by altering the positive potential.³⁹ Using publicly available data, $\epsilon 2$ does not impact expression of nearby genes in GTEx nor does it demonstrate enhancer or promoter chromatin marks in ENCODE HepG2 liver cells supporting $\epsilon 2$'s direct impact on ApoE itself. ApoE $\epsilon 2$ can alternatively clear lipoproteins via cell-surface heparan sulfate proteoglycan and LDL receptor-related protein.⁴⁰⁻⁴² ApoE $\epsilon 2$ transgenic mice crossbred with ApoB transgenic mice have lower LDL cholesterol.⁴² Furthermore, ApoE $\epsilon 2$ transgenic mice lacking LDL receptor still had lower LDL cholesterol suggesting that hypocholesterolemia appears independent of $\epsilon 2$'s effects on LDL receptor.^{35, 43} ApoE $\epsilon 2$ impairs lipoprotein lipase-mediated metabolism of VLDL to LDL potentially through the displacement of ApoCII, an activator of lipoprotein lipase.⁴³ The consequent diminished hepatic cholesterol may subsequently increase LDL receptors for ApoB-containing lipoproteins like LDL.

Interestingly, despite accounting for LDL cholesterol or serum triglycerides, we observe that $\epsilon 2$ still is highly associated with reduced CAC quantity. It is likely that single cross-sectional

measure of lipoproteins, while correlates with, does not fully account for lifelong lipoprotein exposures. ApoE ϵ 2 homozygotes who develop type III hyperlipoproteinemia have a marked increase in remnant lipoprotein particles unlike heterozygotes. Analogously, ApoE ϵ 2-overexpressing mice have increased hepatic VLDL production.⁴² Thus, while ApoE ϵ 2 heterozygotes may have an increase in VLDL production and decreased triglyceride catabolism via lipoprotein lipase, the observation of similar triglyceride levels compared to non-carriers suggests preservation of, or enhanced, clearance of remnant lipoprotein particles. We hypothesize that ApoE ϵ 2's association with reduced subclinical atherosclerosis may be due to increased clearance of both atherogenic LDL and remnant lipoprotein particles through LDL receptor-dependent and -independent pathways. Further work is needed to test this hypothesis.

Our study has several strengths. First, we perform a genetic association meta-analysis across the largest set of individuals to-date for subclinical atherosclerosis in two distinct ancestries. Second, we characterize the association of protein-coding genomic variation, which has not been well studied at the population level, with subclinical atherosclerosis. Third, we explore mechanisms of association through lipoprotein-mediation analyses. Fourth, we provide novel insights with both cross-ethnicity and cross-atherosclerosis trait observations. Fifth, we relate the associations of these subclinical atherosclerosis genetic variants on risk for CHD.

While our study has several strengths, we note some key limitations. First, not all protein-coding variation is catalogued on the exome chip. Due to purifying selection, disruptive protein-coding variation is rare.⁴⁴ By potentially not accounting for the totality of disruptive variation not on the array, variance is increased and power is not optimized for gene-based analyses. Whole exome

sequencing can better address this limitation as such technologies continue to become more cost-effective for large-scale experiments. Second, our analyses of prior associations at non-coding sites are restricted to sites on the exome chip. We were able to robustly replicate prior non-coding association analyses for CAC at 9p21 and 6p24.¹² A prior meta-analysis for CIMT genome-wide association discovered one genome-wide association, an intergenic common variant (rs11781551-A) 385kb from *ZHX2* at 8q24.⁷ No variant with modest linkage disequilibrium with this variant was present on the exome chip thereby limiting ability for replication. An intronic variant in *PINX1* at 8q23 and intergenic variant 2.3kb from *APOC1* at 19q13 previously had suggestive association but no suitable proxies to replicate association were available on the exome chip. Third, our analysis still demonstrates a paucity of genome-wide associations for these quantitative atherosclerotic traits and highlights an important challenge to ongoing CAC association analyses.

Genetic determinants of CHD have been characterized among individuals of European ancestry but the strongest association signals have not replicated in those of African ancestry which may be due to smaller sample sizes hindering statistical power or different key genetic drivers. But now we demonstrate a cardioprotective genetic mechanism in those of European ancestry and African ancestry through the reduction of subclinical atherosclerosis. We propose potential mechanisms and call for renewed attention to *APOE* ϵ 2 in the genesis of atherosclerosis underlying clinical cardiovascular disease. Lastly, given the strong concordance of subclinical atherosclerosis measures and clinical CHD, our findings support a future study of genotypes, subclinical atherosclerosis, and incident CHD.

478 **Sources of Funding**

479 P.N. is supported by the John S. LaDue Memorial Fellowship in Cardiology, Harvard Medical
480 School. M.K. is supported by the NWO VENI grant (VENI, 91616079). Infrastructure for the
481 CHARGE Consortium is supported in part by the National Heart, Lung, and Blood Institute grant
482 R01HL105756. Funding support for the CHARGE Consortium Exome Chip analyses is provided
483 in part by the National Heart, Lung, and Blood Institute grant R01HL120393. Please refer to the
484 Supplementary Materials regarding additional sources of funding.

485 **Disclosures Statement**

486 P.N. reports grant support from Amarin Corporation. I.B.B. is the Executive Director of
487 Analytical Genetics at Regeneron Pharmaceuticals Inc. I.I. reports grant support from Pie
488 Medical Imaging PV. O.H.F. works in ErasmusAGE, which is funded by Nestlé Nutrition
489 (Nestec Ltd.), Metagenics Inc., and AXA.

References

1. Go AS, Mozaffarian D, Roger VL, Benjamin EJ, Berry JD, Blaha MJ, et al. Heart disease and stroke statistics--2014 update: a report from the american heart association. *Circulation*. 2014;129:e28-e292.
2. Libby P, Ridker PM and Hansson GK. Progress and challenges in translating the biology of atherosclerosis. *Nature*. 2011;473:317-325.
3. Nambi V, Chambless L, Folsom AR, He M, Hu Y, Mosley T, et al. Carotid intima-media thickness and presence or absence of plaque improves. *J Am Coll Cardiol*. 2010;55:1600-1607.
4. Polonsky TS, McClelland RL, Jorgensen NW, Bild DE, Burke GL, Guerci AD, et al. Coronary artery calcium score and risk classification for coronary heart disease. *Jama*. 2010;303:1610-1616.
5. Yeboah J, McClelland RL, Polonsky TS, Burke GL, Sibley CT, O'Leary D, et al. Comparison of novel risk markers for improvement in cardiovascular risk. *Jama*. 2012;308:788-795.
6. Baber U, Mehran R, Sartori S, Schoos MM, Sillesen H, Muntendam P, et al. Prevalence, Impact, and Predictive Value of Detecting Subclinical Coronary and Carotid Atherosclerosis in Asymptomatic Adults: The BioImage Study. *J Am Coll Cardiol*. 2015;65:1065-1074.
7. Bis JC, Kavousi M, Franceschini N, Isaacs A, Abecasis GR, Schminke U, et al. Meta-analysis of genome-wide association studies from the CHARGE consortium. *Nat Genet*. 2011;43:940-947.
8. Peyser PA, Bielak LF, Chu JS, Turner ST, Ellsworth DL, Boerwinkle E, et al. Heritability of coronary artery calcium quantity measured by electron beam computed tomography in asymptomatic adults. *Circulation*. 2002;106:304-308.
9. Zhao J, Cheema FA, Bremner JD, Goldberg J, Su S, Snieder H, et al. Heritability of carotid intima-media thickness: a twin study. *Atherosclerosis*. 2008;197:814-820.
10. Goff DC, Jr., Lloyd-Jones DM, Bennett G, Coady S, D'Agostino RB, Gibbons R, et al. 2013 ACC/AHA guideline on the assessment of cardiovascular risk: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation*. 2014;129:S49-73.
11. O'Donnell CJ, Cupples LA, D'Agostino RB, Fox CS, Hoffmann U, Hwang SJ, et al. Genome-wide association study for subclinical atherosclerosis in major arterial. *BMC Med Genet*. 2007;8 Suppl 1:S4.
12. O'Donnell CJ, Kavousi M, Smith AV, Kardina SL, Feitosa MF, Hwang SJ, et al. Genome-wide association study for coronary artery calcification with follow-up in myocardial infarction. *Circulation*. 2011;124:2855-2864.
13. Wojczynski MK, Li M, Bielak LF, Kerr KF, Reiner AP, Wong ND, et al. Genetics of coronary artery calcification among African Americans, a meta-analysis. *BMC Med Genet*. 2013;14:75.
14. Zhang L, Buzkova P, Wassel CL, Roman MJ, North KE, Crawford DC, et al. Lack of associations of ten candidate coronary heart disease risk genetic variants and subclinical atherosclerosis in four US populations: the Population Architecture using Genomics and Epidemiology (PAGE) study. *Atherosclerosis*. 2013;228:390-399.
15. Glazier AM, Nadeau JH and Aitman TJ. Finding genes that underlie complex traits. *Science*. 2002;298:2345-2349.

16. Kiezun A, Garimella K, Do R, Stitzel NO, Neale BM, McLaren PJ, et al. Exome sequencing and the genetic basis of complex traits. *Nat Genet.* 2012;44:623-630.
17. Grove ML, Yu B, Cochran BJ, Haritunians T, Bis JC, Taylor KD, et al. Best practices and joint calling of the HumanExome BeadChip: the CHARGE Consortium. *PLoS One.* 2013;8:e68095.
18. Psaty BM, O'Donnell CJ, Gudnason V, Lunetta KL, Folsom AR, Rotter JJ, et al. Cohorts for Heart and Aging Research in Genomic Epidemiology (CHARGE) Consortium: Design of prospective meta-analyses of genome-wide association studies from 5 cohorts. *Circ Cardiovasc Genet.* 2009;2:73-80.
19. Daniell AL, Wong ND, Friedman JD, Ben-Yosef N, Miranda-Peats R, Hayes SW, et al. Concordance of coronary artery calcium estimates between MDCT and electron beam tomography. *AJR Am J Roentgenol.* 2005;185:1542-1545.
20. Mao SS, Pal RS, McKay CR, Gao YG, Gopal A, Ahmadi N, et al. Comparison of coronary artery calcium scores between electron beam computed tomography and 64-multidetector computed tomographic scanner. *J Comput Assist Tomogr.* 2009;33:175-178.
21. Greenland P, Alpert JS, Beller GA, Benjamin EJ, Budoff MJ, Fayad ZA, et al. 2010 ACCF/AHA guideline for assessment of cardiovascular risk in asymptomatic adults: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol.* 2010;56:e50-103.
22. Agatston AS, Janowitz WR, Hildner FJ, Zusmer NR, Viamonte M, Jr. and Detrano R. Quantification of coronary artery calcium using ultrafast computed tomography. *J Am Coll Cardiol.* 1990;15:827-832.
23. Peloso GM, Auer PL, Bis JC, Voorman A, Morrison AC, Stitzel NO, et al. Association of low-frequency and rare coding-sequence variants with blood lipids and coronary heart disease in 56,000 whites and blacks. *Am J Hum Genet.* 2014;94:223-232.
24. Price AL, Patterson NJ, Plenge RM, Weinblatt ME, Shadick NA and Reich D. Principal components analysis corrects for stratification in genome-wide association studies. *Nat Genet.* 2006;38:904-909.
25. Pruim RJ, Welch RP, Sanna S, Teslovich TM, Chines PS, Gliedt TP, et al. LocusZoom: regional visualization of genome-wide association scan results. *Bioinformatics.* 2010;26:2336-2337.
26. Purcell S, Cherny SS and Sham PC. Genetic Power Calculator: design of linkage and association genetic mapping studies of complex traits. *Bioinformatics.* 2003;19:149-150.
27. Sabatti C, Service SK, Hartikainen AL, Pouta A, Ripatti S, Brodsky J, et al. Genome-wide association analysis of metabolic traits in a birth cohort from a founder population. *Nat Genet.* 2009;41:35-46.
28. Stitzel NO, Stirrups K, Masca NGD, Erdmann J, Ferrario PG, König IR, et al. Coding Variation in ANGPTL4, LPL, and SVEP1 and the Risk of Coronary Disease. *N Engl J Med.* 2016;374:1134-1144.
29. Folsom AR, Kronmal RA, Detrano RC, O'Leary DH, Bild DE, Bluemke DA, et al. Coronary artery calcification compared with carotid intima-media thickness in the prediction of cardiovascular disease incidence: the Multi-Ethnic Study of Atherosclerosis (MESA). *Arch Intern Med.* 2008;168:1333-1339.
30. Mega JL, Stitzel NO, Smith JG, Chasman DI, Caulfield MJ, Devlin JJ, et al. Genetic risk, coronary heart disease events, and the clinical benefit of statin therapy: an analysis of primary and secondary prevention trials. *Lancet.* 2015;385:2264-2271.

31. Innerarity TL, Weisgraber KH, Arnold KS, Mahley RW, Krauss RM, Vega GL, et al. Familial defective apolipoprotein B-100: low density lipoproteins with abnormal receptor binding. *Proc Natl Acad Sci U S A*. 1987;84:6919-6923.
32. Shen H, Damcott CM, Rampersaud E, Pollin TI, Horenstein RB, McArdle PF, et al. Familial defective apolipoprotein B-100 and increased low-density lipoprotein cholesterol and coronary artery calcification in the old order amish. *Arch Intern Med*. 2010;170:1850-1855.
33. Segrest JP, Jones MK, De Loof H and Dashti N. Structure of apolipoprotein B-100 in low density lipoproteins. *J Lipid Res*. 2001;42:1346-1367.
34. Hallman DM, Boerwinkle E, Saha N, Sandholzer C, Menzel HJ, Csazar A, et al. The apolipoprotein E polymorphism: a comparison of allele frequencies and effects in nine populations. *Am J Hum Genet*. 1991;49:338-349.
35. Bennet AM, Di Angelantonio E, Ye Z, Wensley F, Dahlin A, Ahlbom A, et al. Association of apolipoprotein E genotypes with lipid levels and coronary risk. *Jama*. 2007;298:1300-1311.
36. Sudlow C, Martinez Gonzalez NA, Kim J and Clark C. Does apolipoprotein E genotype influence the risk of ischemic stroke, intracerebral hemorrhage, or subarachnoid hemorrhage? Systematic review and meta-analyses of 31 studies among 5961 cases and 17,965 controls. *Stroke*. 2006;37:364-370.
37. Innerarity TL, Pitas RE and Mahley RW. Binding of arginine-rich (E) apoprotein after recombination with phospholipid vesicles to the low density lipoprotein receptors of fibroblasts. *J Biol Chem*. 1979;254:4186-4190.
38. Mahley RW. Apolipoprotein E: cholesterol transport protein with expanding role in cell biology. *Science*. 1988;240:622-630.
39. Dong LM, Parkin S, Trakhanov SD, Rupp B, Simmons T, Arnold KS, et al. Novel mechanism for defective receptor binding of apolipoprotein E2 in type III hyperlipoproteinemia. *Nat Struct Biol*. 1996;3:718-722.
40. Huang Y, Rall SC, Jr. and Mahley RW. Genetic factors precipitating type III hyperlipoproteinemia in hypolipidemic transgenic mice expressing human apolipoprotein E2. *Arterioscler Thromb Vasc Biol*. 1997;17:2817-2824.
41. de Beer F, Hendriks WL, van Vark LC, Kamerling SW, van Dijk KW, Hofker MH, et al. Binding of beta-VLDL to heparan sulfate proteoglycans requires lipoprotein lipase, whereas ApoE only modulates binding affinity. *Arterioscler Thromb Vasc Biol*. 1999;19:633-637.
42. Mahley RW, Huang Y and Rall SC, Jr. Pathogenesis of type III hyperlipoproteinemia (dysbetalipoproteinemia). Questions, quandaries, and paradoxes. *J Lipid Res*. 1999;40:1933-1949.
43. Huang Y, Liu XQ, Rall SC, Jr. and Mahley RW. Apolipoprotein E2 reduces the low density lipoprotein level in transgenic mice by impairing lipoprotein lipase-mediated lipolysis of triglyceride-rich lipoproteins. *J Biol Chem*. 1998;273:17483-17490.
44. MacArthur DG, Balasubramanian S, Frankish A, Huang N, Morris J, Walter K, et al. A systematic survey of loss-of-function variants in human protein-coding genes. *Science*. 2012;335:823-828.

Table 1. Top meta-analysis variant associations for coronary artery calcification quantity

					All				EA		AA	
Variant	Consequence	Nearest Gene*	Chrom:Pos†	Minor Allele	MAF	Beta‡	SE	P	MAF	P	MAF	P
rs10757278 §	intergenic	(<i>CDKN2B</i>)	9:22124477	G	0.43	0.21	0.020	3.14x10 ⁻²⁴	0.48	2.9x10 ⁻²⁵	0.21	0.20
rs9349379	intronic	<i>PHACTR1</i>	6:12903957	G	0.34	0.19	0.020	4.93x10 ⁻²⁰	0.39	1.28x10 ⁻¹⁹	0.094	0.088
rs7412#	missense	<i>APOE</i>	19:45412079	T	0.081	-0.25	0.036	1.19x10 ⁻¹²	0.074	4.43x10 ⁻⁶	0.11	5.36x10 ⁻¹⁰
rs1412829§	intronic	<i>CDKN2B</i>	9:22043926	C	0.34	-0.14	0.021	1.56x10 ⁻¹¹	0.41	5.58x10 ⁻¹²	0.072	0.84
rs5742904**	missense	<i>APOB</i>	2:21229160	T	2.1x10 ⁻³	1.41	0.22	2.93x10 ⁻¹⁰	2.7x10 ⁻³	2.93x10 ⁻¹⁰	0	NA
rs9369640	intronic	<i>PHACTR1</i>	6:12901441	A	0.43	-0.11	0.019	4.91x10 ⁻⁸	0.38	5.04x10 ⁻⁹	0.36	0.71
rs769449#	intronic	<i>APOE</i>	19:45410002	A	0.10	0.14	0.032	7.93x10 ⁻⁶	0.11	1.86x10 ⁻⁶	0.024	0.19
rs1801696**	missense	<i>APOB</i>	2:21232044	T	4.6x10 ⁻³	0.63	0.14	1.44x10 ⁻⁵	5.7x10 ⁻³	9.77x10 ⁻⁶	0	NA

* Genes for SNPs that are outside the transcript boundary of the protein-coding gene are shown in parentheses [eg, (*CDKN2B*)].

† Genomic positions correspond to GRCh37.p13 reference, forward strand.

‡ β-Coefficients are estimated for natural log transformation of total Agatston CAC score+1.

§ *CDKN2B* lead variants show modest correlation among EA (r²=0.24) and no correlation among AA.

CIRCCVG/2016/001572-T

|| *PHACTR1* lead variants show modest correlation among EA ($r^2=0.36$) and AA ($r^2=0.05$)

APOE lead variants show minimal correlation among EA ($r^2=0.01$) and no correlation among AA.

** *APOB* lead variants are not observed to be correlated.

Abbreviations: AA=African ancestry; AF=minor allele frequency; Chrom:Pos=hg19 build chromosome:position; EA=European ancestry; SE=standard error

Table 2. Top meta-analysis variant associations for carotid intima media thickness

Variant	Consequence	Nearest Gene*	Chrom:Pos†	Minor Allele	All				EA		AA	
					MAF	Beta‡	SE	P	MAF	P	MAF	P
rs7412	missense	<i>APOE</i>	19: 45412079	T	0.083	-0.014	0.0022	3.79x10 ⁻¹⁴	0.079	1.97x10 ⁻¹⁰	0.11	1.43x10 ⁻⁵
rs11668477	intergenic	(<i>LDLR</i>)	19:11195030	G	0.27	-0.0064	0.0016	4.69x10 ⁻⁷	0.20	5.26x10 ⁻⁶	0.34	0.030
rs7188	3'UTR	(<i>KANK2</i>)	19:11275139	G	0.29	0.0054	0.0011	2.23x10 ⁻⁶	0.33	1.36x10 ⁻⁶	0.079	0.98
rs1712790	intergenic	(<i>FAM55B</i>)	11:114621469	C	0.47	-0.0048	0.0011	5.93x10 ⁻⁶	0.48	1.89x10 ⁻⁶	0.21	0.89
rs2298375	missense	<i>C22orf15</i>	22:24106448	A	0.086	0.0082	0.0019	9.51x10 ⁻⁶	0.085	5.64x10 ⁻⁶	0.091	0.061
rs174547	intronic	(<i>FADS1</i>)	11:61570783	C	0.30	-0.0049	0.0011	1.07x10 ⁻⁵	0.34	3.84x10 ⁻⁵	0.082	0.062

* Genes for SNPs that are outside the transcript boundary of the protein-coding gene are shown in parentheses [eg, (*LDLR*)].


† Genomic positions correspond to GRCh37.p13 reference, forward strand.

‡ β-Coefficients are estimated for natural log transformation of CIMT.


Abbreviations: AA=African ancestry; AF=minor allele frequency; Chrom:Pos=hg19 build chromosome:position; EA=European ancestry; SE=standard error; UTR=untranslated region

Figures


Figure 1. Association of each genotyped variant with CAC quantity




Plot of $-\log_{10}(P)$ for association of genotyped variants by chromosomal position for all autosomal polymorphisms analyzed in the age-, sex-, and principal components- adjusted model of coronary artery calcification quantity in the meta-analysis. The genes associated with the top associated variants are displayed.

Figure 2. Forest plot of relative CAC quantity for *APOE* ϵ 2 carriers

CAC quantity for *APOE* ϵ 2 carriers relative to non-carriers is displayed for all cohorts stratified by European and African ancestries to demonstrate consistency across diverse cohorts and ethnicities.

Figure 3. Association of each genotyped variant with CIMT

Plot of $-\log_{10}(P)$ for association of genotyped variants by chromosomal position for all autosomal polymorphisms analyzed in the age-, sex-, and principal components- adjusted model of carotid intima media thickness in the meta-analysis. The genes associated with the top associated variants are displayed.

Figure 4. Forest plot of relative CIMT for *APOE* $\epsilon 2$ carriers

CIMT for *APOE* $\epsilon 2$ carriers relative to non-carriers is displayed for all cohorts stratified by European and African ancestries to demonstrate consistency across diverse cohorts and ethnicities.